

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/92498 A1

(51) International Patent Classification⁷: C12N 9/02
// (C12N 9/02, C12R 1:645), D21C 5/00, A61K 7/13,
C11D 3/386

(74) Agent: SEPPO LAINE OY; Itämerenkatu 3 B,
FIN-00180 Helsinki (FI).

(21) International Application Number: PCT/FI01/00503

(22) International Filing Date: 23 May 2001 (23.05.2001)

(25) Filing Language: Finnish

(26) Publication Language: English

(30) Priority Data:
20001240 23 May 2000 (23.05.2000) FI

(71) Applicant (for all designated States except US): VAL-
TION TEKILLINEN TUTKIMUSKESKUS [FI/FI];
Vuorimiehentie 5, FIN-02044 VTT (FI).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE
(utility model), DK, DK (utility model), DM, DZ, EC, EE,
EE (utility model), ES, FI, FI (utility model), GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG,
US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KRUUS, Kristiina
[FI/FI]; Kaitalahdenranta 7, FIN-02260 Espoo (FI). KI-
ISKINEN, Laura-Leena [FI/FI]; Metsätöntunkuja 3 E
19, FIN-02200 Espoo (FI). RÄTTÖ, Marjaana [FI/FI];
Syyrakinkuja 9, FIN-01690 Vantaa (FI). VIIKARI, Liisa
[FI/FI]; Lokkikuja 5 F, FIN-00200 Helsinki (FI). SALO-
HEIMO, Markku [FI/FI]; Kytäjäntie 18 b, FIN-00390
Helsinki (FI).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL LACCASE ENZYME AND THE GENE ENCODING THE ENZYME

(57) Abstract: The invention relates to a new laccase enzyme, which can be isolated from the strains of the *Melanocarpus* genus, the *M. albomyces* strain in particular. The pH optimum of the enzyme is within 5-8 and the enzyme works at a temperature of 30-80 °C. The isoelectric point of the enzyme is about 4.0 as determined by isoelectric focusing and the molecular weight about 80 kDa, defined by SDS-PAGE. The enzyme is especially well suited to applications, wherein the pH and temperature conditions are high. The invention also relates to a gene that encodes laccase enzyme, and a laccase enzyme produced by recombinant technology.



WO 01/92498 A1

Novel laccase enzyme and the gene encoding the enzyme.

The invention relates to new laccase enzymes. The object of the invention is a new laccase enzyme, an enzyme preparation, and a gene encoding the enzyme in particular. Another
5 object of the invention is the use of laccase in various applications.

Laccases (EC 1.10.3.2) belong to blue copper oxidases. According to a definition, laccases are *p*-diphenol oxidases. In addition to diphenols, laccases oxidize many other substrates, such as methoxy substituted phenols and diamines. With respect to their substrates, lac-
10 cases are amazingly unspecific. Namely because of their broad substrate specificity and, on the other hand, their ability to oxidize phenolic compounds, laccases have aroused a great interest in industrial applications. Promising fields of application for laccases include, for example, delignification and the gluing of fibreboards in the forest industry, dyeing of fab-
15 rics and detoxification of dye house effluents in the textile industry, as well as use in different biosensors. With the aid of mediators, i.e., intermediary molecules, laccases also oxidize substrates that they otherwise would not be able to oxidize. The mediators are small-molecule compounds that are oxidized by laccases. The oxidized mediator, in turn, oxidizes the actual substrate.

20 The first laccase was found as early as 1883 in the Japanese lacquer tree (*Rhus vernicifera*). Laccases have been found in many plants, such as peach, tomato, mango, and potato; laccases have also been found in some insects. However, most known laccases originate in white rot fungi. The following genera, for example, produce laccase: *Agaricus*, *Aspergillus*, *Cerrena*, *Curvularia*, *Fusarium*, *Lentinius*, *Monocillium*, *Myceliophthora*, *Neurospora*,
25 *Penicillium*, *Phanerochaete*, *Phlebia*, *Pleurotus*, *Podospora*, *Schizophyllum*, *Sporotrichum*, *Stagonospora* and *Trametes*. In nature, the functioning of laccases relates to the decomposition of lignocellulose, the biosynthesis of cell walls, the browning reactions of fruit and vegetables, as well as the prevention of microbial attacks on plants, among others.

30 Many fungal laccases have been isolated and some genes that encode them have been cloned. For example, Saloheimo *et al.* (1985) isolated and characterized the laccase gene of

the *Phlebia radiata*, and Kojima *et al.* (1990) the laccase genes of the *Coriolus hirsutus* from white-rot fungus, and Berka *et al.* (1997), WO 95/33836, the gene of the *Myceliophthora thermophila* laccase. The natural production levels of laccases are often very low. Efforts have been made to improve production by expressing laccase genes in foreign production hosts. For example, Saloheimo and Niku-Paavola (1991; WO92/01046) successfully produced *Phlebia radiata* laccase in the fungus *Trichoderma reesei*. Laccases have also been produced heterologously, in the *Aspergillus oryzae* fungus (Yaver *et al.*, 1996, Berka *et al.* 1997 and WO 95/33836), and in the *Pichia pastoris* yeast (Jönsson *et al.*, 1997).

The expression of a laccase originating in the *Coprinus* genus in the fungus of the *Aspergillus* genus is described in the patent publication WO 97/08325. Similarly, the expression of the laccase originating in the *Polyporus pinsitus* species and the laccase originating in the *Scytalidium* genus in the fungus of the *Aspergillus* genus are described in the patent publications US 5,770,418 and US 5,843,745, respectively.

The temperature and pH properties of laccases isolated from various organisms differ from each other. They also depend on the substrate used. As the majority of laccases that are known function best at an acidic pH and a rather low temperature, their properties are not optimal for the applications. Some thermostable or neutral laccases have been reported but, generally, the thermostable or neutral laccases are either thermostable or neutral, not both. For example, Heinzkill *et al.* (1998) found laccases with an unusually high pH optimum from the fungi of the *Coprinaceae* genus. However, in 60 minutes, the activity of all the laccases that were found decreased to below 30% from the initial level at 60°C. A patent (WO96/06930) has been applied for the use of these laccases in bleaching textile dyes, but the application does not mention the activity of these laccases at high temperatures. The application examples were made at a temperature of 30-35°C. Patent application WO95/07988 describes a neutral laccase from the fungus *Rhizoctonia solani*, but the application does not study the applicability of this laccase at high temperatures in addition to alkaline conditions. On the other hand, the patent application WO98/55628 describes a thermostable laccase from the *Trametes versicolor* TV-1 strain, but according to the patent,

the activity of this laccase is best at pH 2. The *Trametes versicolor* TV-1 laccase is not pH stable either; its residual activity at pH 6 after 60 minutes of incubation is about 60% of the original. Patent application WO95/33836 describes a neutral *Myceliophthora thermophila* laccase that in hair dyeing works at pH 7, but the temperature in the application example
5 was 30°C. The publication Berka et al. (1997) reports that the *Myceliophthora thermophila* laccase retains 100% of its activity at 60°C for 20 minutes and its optimum activity is at pH 6.5.

Patent publications JP8070861 and JP9056378 describe *Trametes* laccases, which have
10 been reported to be thermostable and their pH optimum has been reported to be 5.0. In addition, Diamantidis et al. (2000) have characterized a bacterial laccase from the *Azospirillum lipoferum* bacterium. They reported that the laccase was thermostable for 10 min at 70°C and the pH optimum was 6.0.

15 The publication Bharathi and Ramalingam (1993) describes the phenoloxidase of a clam, its maximum activity being at pH 6.8 and more than 50% of its activity remaining when incubated at 60°C for 10 min. Patent publication EP-A1-0852260 describes polyphenoloxi-
dases from the species of the *Myrothecium* genus, their pH optimum being 8.5-9 and the optimum temperature at 60-70°C. It is known that the fungi of the *Myrothecium* genus pro-
20 duce toxins.

When comparing the temperature and the pH optimum values specified for various en-
zymes, it should be noted that the substrate used has an effect on the values. Phenolic sub-
strates, such as guaiacol and syringaldazine, provide higher values than non-phenolic sub-
25 strates, such as ABTS.

The following patent publications suggest the use of laccases in wood-processing applica-
tions: WO 9954545, US 5,691,193, DE 4137761, EP 408803 and WO 9523232.

30 The purpose of this invention is to eliminate the disadvantages related to prior art and to provide quite a new kind of enzyme preparation containing laccase activity. Especially, the

enzyme preparation can be used in applications, which need to tolerate both high temperatures and a high pH. There are such applications, for example, in the wood-processing industry and other processing industries.

- 5 The present invention introduces a new laccase that has both better temperature and pH properties, and especially the thermal stability, than the known laccases in applications that require a neutral or alkaline pH and a temperature of over 40°C.

To be more precise, the object of the invention is a laccase enzyme specified in claim 1.

10

The laccase enzyme according to the invention can be isolated from the strains of the *Melanocarpus* genus. As far as we know, no laccase isolated from the strains of the *Melanocarpus* genus has been described earlier. The laccase according to the invention can preferably be isolated from the strains of the *M. albomyces* species, especially from the IMI
15 255989 strain, samples of which are freely available from the CABI Bioscience UK centre CABI GRC strain collection (Egham) (Bakeham Lane Egham Surrey DW20 9Ty, UK). The strain was deposited in the collection in 1981. In 1996, the strain was also deposited in the Culture Collection of VTT Biotechnology (Technical Research Centre of Finland VTT, at the address: VTT Biotekniikka, PL1500, 02044 VTT, Espoo, Finland) and given the
20 number VTT-D-96490. Samples of this deposition are also freely available. This strain is described in the publication Ravanko (1996), which studied the laccase activity of the culture solution of the strain at various temperatures and pH values. However, the publication does not describe the isolation and the purification of the laccase.

- 25 The laccase enzyme according to the invention works over the range of pH 3-9, preferably 4-8, more preferably 5-8, even more preferably 6-8, most preferably 6.5-7.5. The laccase activity is at its highest over the range of pH 7-8, the highest at about 7.5. Thus, the pH optimum of the enzyme is quite wide and ranges over pH 5-7.5; preferably the pH optimum is 7.5. The activity of the enzyme is at its highest within 30-80°C, preferably within 40-80°C,
30 more preferably within 50-80°C, even more preferably within 60-80°C, most preferably within 60-70°C. The best activity of the enzyme occurs at a temperature of about 70°C.

The laccase according to the invention retains well its activity at high temperatures. In the conditions described in Example 3, more than 50% of the activity of the laccase of the culture solution remains after incubating at 60°C for 1 hour. After incubating at 70°C for 15 min, about 30% of the activity of the enzyme remains, and after incubating at 70°C for 30 min, about 10%. At 80°C, the enzyme withstands incubation for about 5 min in the conditions described in Example 3. In some applications, e.g., in the forest industry, even a tolerance for a short time at high temperatures provides considerable advantages. When further studying the activity of the pure enzyme, it was discovered that the thermal stability of the enzyme was exceptionally good and clearly better than that of the laccases previously described. At 60°C, the enzyme retained its activity essentially unchanged for 2 hours, as indicated by Example 3 and Fig. 7. The thermal stability of the *Myceliophthora* laccase characterized by Berka et al. (1997) remained at 60°C for 20 min only.

Furthermore, the pH stability of the laccase enzyme according to the invention improves, when the pH increases. After 22-hour incubation at pH 4, the residual activity is 65% , the residual activity in the same conditions at pH 5 is about 80%, the residual activity at pH 6 is about 85%, the residual activity at pH 7 is about 90%, and at pH 8 as much as over 90%, about 92%. Together with the pH optimum, this property makes it advantageous to use the enzyme according to the invention at high pH values.

The isoelectric point of the laccase according to the invention is about 4.0 when measured by isoelectric focusing, whereby the accuracy of the definition is about ± 0.5 .

When determined by SDS-PAGE, the molecular weight of the laccase according to the invention is about 80 kDa. The accuracy of the SDS-PAGE definition is in the order of ± 5 kDa.

The objective of the invention is a laccase in particular, which, when purified, has an isoelectric point of about 4.0 as measured by isoelectric focusing, and a molecular weight of about 80 kDa. The pH optimum of the laccase is about 7.5 and the temperature optimum about 70°C when measured with a purified enzyme. The laccase can be isolated from the

strains of the *Melanocarpus* genus, preferably from the strains of the *Melanocarpus albomyces* genus, most preferably from the *M. albomyces* IMI 25598 strain. Another objective of the invention is an isolated and purified laccase enzyme.

- 5 A further object of the invention is an enzyme preparation containing the laccase enzyme according to the invention. The amount of laccase enzyme in the enzyme preparation is preferably higher than that of the strains of the *Melanocarpus* genus, especially the strains of the *Melanocarpus albomyces* species, especially the amount of enzyme naturally produced by the *M. albomyces* IMI 255989 strain in its culture solution in growth conditions,
10 which are not optimised to produce laccase. The amount of laccase in the enzyme preparation is preferably over 10 mg/l, more preferably ≥ 30 mg/l, even more preferably ≥ 300 mg/l, still even more preferably ≥ 500 mg/l; most preferably ≥ 1 g/l.

- As stated above, the laccase according to the invention can be isolated from the strains of
15 the *Melanocarpus* genus, especially from the strains of the *Melanocarpus albomyces* species, especially from the *M. albomyces* IMI 255989 strain, but it can also be produced by the recombination technique by isolating genes that encode the laccases according to the invention and transferring them into a suitable production host.

- 20 The laccase according to the invention can be produced in the culture solution of its natural host or a production host, from where it can be isolated and purified by using known methods of protein chemistry. If the culture solution contains a high enough amount of laccase, but no other detrimental proteins, the culture solution may be used as such by simply separating the cells. The culture solution can possibly be concentrated. In different applica-
25 tions, it is preferable to use an enzyme preparation containing an increased amount of laccase. The increased amount of laccase can be prepared by producing the laccase enzyme in the culture solution of the production host by means of recombinant technology. The increased amount refers to an amount of laccase enzyme, which exceeds the amount of laccase enzyme naturally produced by the strains of the *Melanocarpus* genus, especially the
30 *M. albomyces* strain, especially the IMI 255989 strain.

A further object of the invention is an enzyme preparation containing an essential amount of the laccase enzyme according to the invention. This means that the laccase is the main activity of the enzyme preparation without any considerable amounts of other enzymes. A further object of the invention is also an enzyme preparation containing a laccase that can be isolated from the fungus of the *Melanocarpus* genus, and additives that are needed for the respective application. Such additives can comprise, for example, buffers and stabilizers.

An object of the invention is also a nucleic acid molecule that encodes laccase. The nucleic acid molecule is selected from the group comprising:

- a nucleic acid molecule comprising the coding region of the nucleotide sequence as depicted in SEQ ID NO:1 or Fig. 15;
- a nucleic acid molecule that encodes a polypeptide that comprises the amino acid sequence as depicted in SEQ ID NO:2 or Fig. 15;
- a nucleic acid molecule comprising a coding sequence that differs from the coding sequence of the nucleotide sequence of SEQ ID NO:1 or Fig. 15 due to the degeneracy of the genetic code;
- a nucleic acid molecule that hybridizes to the nucleotide sequence of SEQ ID NO: 1 or Fig. 15; and
- a nucleic acid molecule that encodes a polypeptide, which has laccase activity and the amino acid sequence of which shows at least 73% identity with the amino acid sequence of SEQ ID NO:2 or Fig. 15.

The invention also embodies nucleic acid molecules that encode polypeptides, the amino acid sequence of which shows at least 75% identity with SEQ ID NO:2 or the amino acid sequence of Fig. 15, preferably at least 80% identity, more preferably at least 85% identity, even more preferably at least 90% identity; most preferably at least 95% identity. It is most preferable if the amino acid sequence of the polypeptides according to the invention shows at least 99% identity with ID NO:2 or the amino acid sequence of Fig. 15 . If the similarity is assessed on the basis of homologous amino acids, the invention embodies the nucleic acid molecules, which encode polypeptides, the amino acid sequence of which is at least

83% homologous to SEQ ID NO:2 or the amino acid sequence of Fig. 15, preferably at least 85% homologous, more preferably at least 90% homologous, even more preferably at least 95%, most preferably at least 99% homologous to SEQ ID NO:2 or the amino acid sequence of Fig. 15.

5

The invention also embodies the nucleic acid molecules, which are hybridised to the sequence SEQ ID NO:1 in hybridising conditions, where the hybridisation solution contains 6XSSC, 5XDenhardt's reagent, 0.5% SDS, 100 µg/ml denatured DNA and hybridisation is carried out at 50-60°C. Another alternative is that the hybridisation solution contains

10 6XSSC, 5XDenhardt's reagent, 0.5% SDS, 100 µg/ml denatured DNA, 50% formamide and, in that case, hybridisation is carried out at 25-35°C. The invention especially embodies those nucleic acid molecules, which are hybridised to the SEQ ID NO:1 sequence in stringent conditions, whereby hybridisation is carried out in accordance with the first alternative, otherwise in the same conditions, but at 68°C and, according to the second
15 alternative, at 42°C. 50x Denhardt is 10 g/l Ficoll, 10 g/l polyvinyl pyrrolidone, 10 g/l bovine serum albumin and the SSC is 0.15 M NaCl, 0.015 M sodium citrate, pH 7.0

The nucleic acid molecules refer to DNA, RNA or, for example, cDNA.

20 The invention also relates to polypeptides, which have laccase activity and which are encoded by the nucleic acid molecules defined above. The invention further relates to polypeptides, the amino acid sequence of which, in the way described above, is identical or homologous with SEQ ID NO:2 or the amino acid sequence of Fig. 15.

25 The invention also relates to a method for producing laccase, comprising the following steps:

- a nucleic acid molecule or vector according to the invention is transferred into a micro-organism host cell to express the nucleic acid molecule and, optionally, to secrete it from the host cell; and

- a polypeptide that has laccase activity is recovered either from the cells or the culture solution of the micro-organism host.

The laccase according to the invention is exceptionally well suited to industrial applications, wherein the prevailing pH and temperature conditions are high. Such applications include those of the forest industry, which affect lignin or extractives (either directly or by mediators), manufacture of fibre products and boards from mechanically ground lignin-containing fibres, improvement of the runnability of paper machines, and other applications, oxidation of polymers, such as lignin, cellulose, and/or starch (either directly or by mediators), as well as oxidation of other chemicals, such as alkenes or colour molecules. In these applications, the temperature is often over 60°C, generally as high as 80°C, and the pH is close to neutral or slightly higher. The enzyme according to the invention works extremely well in conditions, where both the pH and the temperature are simultaneously rather high.

In the following, the invention is described in detail with the aid of the appended figures and examples.

Fig. 1 shows the production of the *M. albomyces* laccase in shake flask cultivation.

Fig. 2 shows the laccase activity and the protein content of various fractions in anion exchange chromatography.

Fig. 3 shows the laccase activity and the protein content of various fractions in hydrophobic interaction chromatography.

Fig. 4 shows the dependence of *M. albomyces* laccase activity on the temperature.

Fig. 5 shows the activity of the *M. albomyces* laccase at various pH values.

Fig. 6 shows the residual activity of the *M. albomyces* laccase at various temperatures.

Fig. 7 shows the residual activity of a pure *M. albomyces* laccase at various temperatures.

Fig. 8 shows the residual activity of the *M. albomyces* laccase at various pH values after 22-hour incubation.

Fig. 9 shows the ability of *M. albomyces* and *T. hirsuta* laccases to form colour from 2,6-dimethoxyphenol at 40°C at various pH values. The amount of laccase was 15 nkat/mmol of substrate.

Fig. 10 shows the ability of the *M. albomyces* and *T. hirsuta* laccases to form colour from 2,6-dimethoxyphenol at 60°C at various pH values. The amount of laccase was 30 nkat/mmol of substrate.

Fig. 11 shows the decolourisation of a textile dye under the effect of the *Melanocarpus* and
5 *Trametes* laccases.

Fig. 12 shows the polymerisation of a lignin model substance under the effect of the *Melanocarpus* and *Trametes* laccases at 40°C.

Fig. 13 shows the polymerisation of the lignin model substance under the effect of the *Melanocarpus* and *Trametes* laccases at 60°C.

10 Fig. 14 shows the polymerisation of soluble compounds originating in lignin and extrac-
tives under the effect of the *Melanocarpus* and *Trametes* laccases at 70°C.

Fig. 15 shows a gene that encodes the *Melanocarpus albomyces* laccase, and a corre-
sponding amino acid sequence.

15 The term “enzyme preparation” used in this application refers to any product that contains
laccase enzyme activity. The enzyme preparation can be, for example, a culture solution
containing laccase, an isolated laccase or an enzyme mixture, at least one component of
which is a laccase. The enzyme preparation can also contain various additives, such as
stabilizers or buffers. They are selected so as to be suitable for the respective application of
20 the laccase enzyme preparation. The enzyme preparation can also contain other enzyme
activities, such as peroxidase activity, depending on the application of the enzyme prepara-
tion.

The enzyme preparation containing laccase can also contain a suitable mediator, which is
25 used to enhance the operation of the laccase. Suitable mediators include, e.g., Tempo (= 2,2,6,6-tetramethyl-1-piperidinyloxy), HBT(=1-hydroxybenzotriazole), ABTS=2,2'-
azinobis-3-ethylbenzthiazole-6-sulphonate, violuric acid, NHA (=N-hydroxy-acetanilide).

Screening of microbes that produce laccase

Microbes that produce laccase can be isolated from nature or they can be screened from already isolated and identified strains of culture collections by using screening methods that are well known in microbiology. As laccases belong to phenoloxidases, methods that are suitable for screening phenoloxidases are used in the screenings. Screenings can be carried out studying the production of the enzyme either on a solid culture on plate cultivations or in a liquid culture medium by measuring the enzyme activity.

When looking for new laccases that tolerate higher temperatures and pHs than the majority of known laccases, it is worthwhile to screen microbes from environments, where they live in warm and/or alkaline conditions. Such environments are found, for example, in composts, piles of wood chips or tropical areas.

The production of phenoloxidases by the positive fungi found in the screening can be studied on plates by adding substrates of phenoloxidases on top of the mycelium. By using these drop tests, it can be found out, whether the positive reactions on the plates are caused by peroxidases or laccases. Suitable reagents include ABTS, syringaldazine, and guaiacol, and in observing peroxidases, hydrogen peroxide.

20

The micro-organism, which produces phenoloxidase and which is found as a result of screening is cultivated on a suitable medium, and the formation of phenoloxidase in the culture solution is observed by a method that is suitable for measuring the phenoloxidase activity. Suitable culture media for fungi include, for example, malt extract and potato dextrose media, and suitable substrates for measuring the activity include ABTS, guaiacol, and syringaldazine. For many fungi, the production of phenoloxidases requires an inducer. These include, for example, aromatic compounds, materials containing lignin, surface-active agents, certain sources of carbon, and copper sulphate. The laccase activity can be measured by using ABTS as a substrate, the laccase oxidizes it dark green. The measurement can be made in accordance with the method of Niku-Paavola *et al.* (1988). The lac-

30

case activity can also be measured by using a guaiacol method in accordance with the method of Paszczynski *et al.* (1985).

After a sufficient amount of an interesting laccase has been produced, the enzyme is purified and its properties are characterised. The temperature and pH behaviour as well as the isoelectric point of the enzyme can be determined.

Laccase producers can also be screened by the homology of the sequences of the laccase genes. In that case, nucleotides based on the conserved regions of the amino end of the laccase genes can be used as primers in PCR, and look for gene sequences, which are homologous to known laccases, in the genome of various fungi, for example.

Determination of laccase activity at various temperatures can be carried out by the ABTS method, as described in Example 1. The pH optimum of the laccase can be determined by the guaiacol method in a suitable buffer at different pH values. When the pH is over 7, the operation of the ABTS in the activity analysis weakens.

The thermal stability can be determined by incubating an enzyme sample at various temperatures in a suitable buffer at a certain pH. The residual activity of the enzyme at each temperature can be defined by the ABTS method, for example.

The pH stability can be determined by incubating the enzyme sample at various pH values in a suitable buffer. The residual activity can be determined by the ABTS method, for example.

Isolation and purification of laccase

The enzyme can be purified by using conventional methods of enzyme chemistry, such as salt precipitation, ultrafiltration, ion exchange chromatography, and hydrophobic interaction chromatography. Purification can be monitored by SDS polyacrylamide gel electrophoresis. The enzyme activity of the purified enzyme at various temperatures and pH val-

ues can be determined; similarly, the molecular weight and the isoelectric point can be determined.

In the examples of this invention, the production of laccase is described by cultivating the *Melanocarpus albomyces* strain on a rich culture medium. During cultivation, the laccase

5 activity was monitored in accordance with the ABTS method of Niku-Paavola *et al.* (1988), based on the generation of a dark green cation radical from the ABTS under the effect of the laccase. However, the invention is not limited to the *M. albomyces* laccases produced on this medium only.

10 The *M. albomyces* laccase was purified from a culture solution, from which the cells were removed by filtration or centrifugation. The culture solution was further concentrated using ultrafiltration. An anion exchange chromatography, and hydrophobic interaction chroma-
tography were used for further purification. The molecular weight cut off value in ultrafil-
tration was about 30 kDa. The anion exchange chromatography was carried out at pH 5 in
15 an acetate buffer, and the laccase was eluted from the column with an increasing linear gradient of Na₂SO₄. The best fractions of the ion exchange were further purified by hydro-
phobic interaction chromatography, at pH 5 in an acetate buffer. The sample was bound at 0.7 M Na₂SO₄ concentration and eluted with a decreasing linear gradient of Na₂SO₄.

20 However, it is possible to separate the enzyme by using other known purification methods as well.

The molecular size, the isoelectric point, and the pH and temperature profiles of the purified *M. albomyces* laccase were determined.

25

The purified enzyme refers to an enzyme preparation, which has no other proteins in addition to the laccase band, which can be observed, as defined by SDS-PAGE and Coomassie staining. In this application, the enzyme is purified by ultrafiltration, anion exchange chromatography, and hydrophobic interaction chromatography. The purity of the obtained lac-
30 case that is essentially free from other proteins is ≥90%.

The molecular size of the purified *M. albomyces* laccase was 80 kDa, determined by SDS polyacrylamide gel electrophoresis. In isoelectric focusing by the Multiphor II electrophoresis equipment (Pharmacia LKB), 4.0 was determined as the isoelectric point. The activity of the *M. albomyces* laccase was highest at the temperature of 70°C and its pH optimum was 7.5. After 2 hours of incubation at 60°C, the *M. albomyces* laccase retained 100 % of its activity. Furthermore, its pH stability improved, when the pH increased.

Production of laccase

The laccase according to the invention can be produced in the culture solution of its natural host or a production host, from where it can be isolated and purified by using known methods of protein chemistry. If the culture solution contains a sufficiently high amount of laccase but no other detrimental proteins, it may be possible to use the culture solution as such by simply separating the cells. When so desired, the culture solution can be concentrated and/or purified. It is preferable to use, in various applications, an enzyme preparation containing an increased amount of laccase. Such an enzyme preparation can be prepared by producing the increased amount of laccase enzyme in the culture solution of the production host by means of gene technology. The increased amount refers to an amount of laccase enzyme, which exceeds the amount of laccase enzyme naturally produced by the *M. albomyces* strain, strain IMI 255989 in particular.

The laccase according to the invention can also be produced by recombinant technique by isolating the genes that encode the laccases according to the invention and by transferring them to a suitable production host. The laccase-encoding genes can be isolated from a cDNA library in any of the following ways. The cDNA library can be built in a suitable yeast expression vector and transformed into *Saccharomyces cerevisiae* yeast, for example. The clones that produce laccase are identified, for example, on the basis of activity by using plates that contain ABTS substrate. Another possibility is to connect the cDNA library to a λ ZAP vector and infect the *Escherichia coli* cells with the obtained production bank. The clones that encode laccase are identified either with the aid of a polyclonal antibody or DNA hybridisation. If polyclonal antibodies are used, they are produced against a

purified laccase protein in a rabbit, for example. In hybridisation, fragments of the laccase gene provided by PCR are used as a probe. In that case, the sequences of the PCR primers are either based on the regions generally conserved in the laccase genes (e.g., regions that correspond to the amino acids of the active centre) and/or on the sequence of the amino-terminal end of the purified laccase protein or that of an internal peptide. Furthermore, the oligodT region bound to the polyA region of the tail of each messenger RNA can be used as the PCR primer.

The isolated laccase cDNAs are sequenced. The connection between the isolated cDNA and the isolated enzyme can be ascertained by amino acid sequencing, which is made of the enzyme. The chromosomal copy of the laccase gene can either be isolated by PCR or from the genomic library made in the λ vector, and the locations of the introns can be established by sequencing.

15 Isolation of the gene of the *M. albomyces* laccase

It was not possible to isolate the gene of the *M. albomyces* laccase on the basis of the messenger RNA and the cDNA library. Creating a representative cDNA library was not successful. Surprisingly, a problem was caused by the fact that *M. albomyces* produces laccase in the late cultivation phase, when many cells were already autolyzed, and the messenger RNA partially degraded or it is difficult to isolate pure. Therefore, instead of the cDNA library, the gene had to be isolated from a genomic gene library. The isolation of the gene is described in detail in Example 9.

25 The sequence of the laccase gene is shown in Fig. 15 and in the sequence listing (SEQ ID NO:1). The length of the gene is 2279 bp, including introns. As Fig. 15 shows, the number of the base that starts the encoding region is 286, the introns are in positions 541-618, 698-770, 783-869, 1913-1999, 2069-2150. The last encoding base is 2561.

30 The gene encodes a polypeptide of a length of 623 amino acids. When comparing the amino acid sequence, by using the Blast method (Altschul et al., 1990), with the amino acid

sequence of *Myceliophthora* disclosed in patent application WO 9533836, it was discovered that the amino acid sequences were 72% identical (identities 450/623, 72%) and 82% homologous (positives 518/623, 82%; positives refer to homologous amino acids), (gaps=4/623, 0%).

5

When a corresponding comparison was made with *Podospora anserina* laccase (Fernandez-Larrea and Stahl, 1996), it was discovered that the sequences were 68% identical (identities 427/627, 68%), and 79% homologous (positives 502/627, 79%) (gaps=12/627, 1%).

10

The isolated laccase gene is utilized in protein production in other organisms. Such production hosts include the *Aspergillus* production systems mentioned above, such as *A. oryzae* or *A. niger* (US 5,843,745, US 5,770,418, WO 9708325 and WO 9533386); the production system developed for the fungus *Trichoderma* (EP 244 234), or the production system developed for fungal species of *Fusarium*, such as *F. oxysporum* (Malardier et al., 1989), the production systems developed for a *Bacillus* bacterium, such as the *B. subtilis* or the *E. coli* bacterium, the yeasts *Saccharomyces*, *Shizosaccharomyces* or *Pichia pastoris*, or a *Streptomyces* actinomycete or some other microbe or mammalian cell.

15

20 **Optimizing the production of laccase**

The production of laccase can also be improved by optimising the culture conditions and the culture medium of a wild or a recombinant strain. When optimising the culture medium, for example, the effect of the quality (among others, an organic or inorganic source of nitrogen) and the quantity of the source of nitrogen on the laccase production. When needed, the source of nitrogen is limited in order to achieve a higher yield. Similarly, the effect of the source of carbon is established. A source of carbon that is optimal for the enzyme production is selected. When needed, the amount of carbon source can also be limited. The carbon/nitrogen ratio is optimised to be the best for the production of enzyme.

The growing conditions are optimised to be the best possible for the enzyme production in question. The microbe is grown at a pH and temperature optimal for enzyme production.

25

30

Proper mixing and air supply guarantee an optimal aeration during fermentation. In fermentation, inducers of laccase production, such as veratryl alcohol, xyloidine, or lignin can also be used. The way and the time of adding the inducers, as well as their concentration are optimised.

5

Use of laccase in various applications

Generally, the laccase according to the invention is well suited to be used in applications, wherein laccases can be used, such as gel formation, gluing of fibres, treatment of cork,
10 removal of dye (especially textile dyes), dyeing of fibres, protein treatment, detergents, anti-microbial applications, starch applications, oxidation of chemicals, removal of biofilm, preparation of lignin derivatives, medical analysis, reducing the shrinkage of wool, baking, improving the preservability of beer, production of dye, removal of oxygen from oil products, and the production of iodine. When so desired, the laccase according to the
15 invention can also be immobilized for certain purposes.

The laccase according to the invention is especially well suited to industrial applications, wherein the prevailing pH and temperature conditions are high. Such applications include, among others, applications of the forest industry, manufacture of fibre products and boards
20 from mechanically ground, lignin-containing fibres, improvement of the runnability of paper machines and other applications, oxidation of polymers and that of other chemicals, such as dye molecules. In these applications, the temperature is often over 60°C, generally as much as 80°C, and the pH near neutral or slightly alkalic.

25 In all applications, it is essential that enough oxygen is brought to the reaction. Oxygen is the reducing substrate needed by the laccase. In some cases, especially when the substrate concentration is low, there can be enough oxygen in the reaction mixture as such but it can also be added either by bringing air or oxygen or air enriched with oxygen to the reaction mixture.

30

When needed, mediators are used as additives in the reactions.

The laccase according to the invention is well suited to oxidation of colouring agents, for example. The dye molecules are brought into contact with the laccase at pH 4-8, for example, at pH 4.5-7.5, preferably at pH 6-8, more preferably at pH 7-8, the temperature being within a range of 25-80°C, preferably 40-80°C, more preferably 50-80°C; most preferably 60-80°C. Oxygen is added into the reaction as needed. The amount of laccase is 1-1000 nkat/g, preferably 10-500 nkat/g, most preferably 20-200 nkat/g of colouring agent. The reaction is allowed to take place for 5 min to 24 hours, preferably for 30 min to 2 hours.

The laccase according to the invention is also well suited to polymerisation. The selected compound to be polymerised is brought into contact with the laccase in the same conditions as described above. The reaction mixture can be aerated during the test. Polymerisation can be monitored by following the increase in the molecular weight of the polymerised compound, for example by GPC (gel permeation chromatography).

The laccase according to the invention can also be used to improve the runnability of paper machines. The laccase can be used to improve the runnability of paper machines by polymerising compounds originating from lignin and extractives and by decreasing the detrimental growth of microbes in the paper machine. Generally, the conditions in the paper machine are about pH 5-7 and the temperature 60-80°C. The laccase can be added into the process water or into the head box or the circulation water system of the paper machine without essentially needing to change the conditions prevailing at the paper machine. The pH can be within 5-8 and the temperature within 50-80°C. The amount of laccase can be 1-1000 nkat/g, preferably 10-500 nkat/g of dry matter, fibres or litres of circulation water. The treatment time can be 5 min to 24 hours, preferably 30 min to 2 hours.

The laccase according to the invention can also be used in oxidation of fibres. Lignin-containing fibres can be brought into contact with the laccase at a temperature of 50-80°C, even as high as 50-100°C, preferably at a temperature of 60-80°C, at pH 5-8, preferably at

pH 6-8, the laccase concentration being 1-1000 nkat/g of fibres, preferably 10-500 nkat/g of fibres, the reaction time being 2 min to 24 hours, preferably 10 min to 2 hours. Due to the laccase treatment, the strength properties of the fibres improve, which can be utilised, for example, in the manufacture of fibre boards, such as MDF boards, or in paper or card-board products, which are made of mechanically ground lignin-containing fibres.

In addition to the application examples mentioned above, the laccase according to the invention can also be used in delignification of fibres. The laccase can be in contact with the fibres that are to be delignified, such as kraft fibres, with the kappa value being 20-30, preferably about 25, the consistency being 5-15, preferably about 10 %, preferably in the presence of a mediator the amount of mediator being 1-5% of the pulp, preferably 3% of the pulp, the pH being within 5-8, preferably within 6-8, the temperature being within 50-80°C, preferably within 60-80°C. The reaction time can be between 5 min to 24 hours, preferably 30 min to 2 hours. The amount of laccase can be 10-1000 nkat/g, preferably 10-500 nkat/g in an oxygen pressure of 0.5 Mpa.

The following examples are intended for illustration of the present invention and should not be interpreted as limiting the present invention in any way.

Example 1.

Production of *M. albomyces* laccase

M. albomyces fungus was kept on oat-agar plates (Difco). Both the inoculum and the production medium contained

25 g/l glucose (AnalaR)

27.5 g/l Bacto yeast extract (Difco)

0.5 mg/ml Indulin AT (Sigma)

0.04 l/l mineral solution containing:

1.0 g/l $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (Riedel-de Haën)

1.0 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (Riedel-de Haën)

0.1 g/l $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (Merck)

0.16 g/l $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (Merck)
1.0 g/l Na_2EDTA (Riedel-de Haën).

The glucose solution was sterilized separately.

5

First, 100 ml of medium was inoculated with 3-4 pieces (about 1 cm^2) cut from well grown mycelium on oatmeal agar. The cultivation temperature was 37°C and the agitation speed 120 rpm. After 2 days of cultivation, the mycelium was homogenized, and 900 ml of the sterile culture was inoculated with 100 ml of the homogenized inoculum. The volume of the production culture was 1 l; the cultivation temperature was 37°C , and the agitation speed 160 rpm. Cultivation was continued for 14 days. Four parallel cultures were made.

Enzyme activity assay

The laccase activity of the *M. albomyces* culture solutions was measured by using ABTS. Laccase oxidizes ABTS to a dark green cation radical. The activity assay was carried out in accordance with the method developed by Niku-Paavola *et al.* (1988). The sample was diluted with 0.025 M succinate buffer, pH 4.5. 0.350 ml of ABTS solution (11 g/l) was added into 1.15 ml of the dilution, and the reaction was followed for 2 minutes by the Perkin Elmer Lambda 20 spectrophotometer at a wavelength of 436 nm.

20

The measured laccase activities of the *M. albomyces* cultures are shown in Fig. 1.

Example 2.

Purification of *M. albomyces* laccase

25

Determination of the protein content

The protein contents were determined by the DC Protein Assay kit of Bio-Rad, based on a method developed by Lowry *et al.* (1951). The assays were carried out using the reagents of the kit, and the intensity of the colour reaction thus formed was measured on a wavelength of 750 nm by the Hitachi U-2000 spectrophotometer. Each time of measurement, a

30

standard curve was also defined using solutions containing 0.25-1.25 g/l of bovine serum albumin (BSA, Bio-Rad).

Purification

5 The culture solution, from which cells had been removed by filtration, was ultrafiltrated by the Amicon 8400 filtering equipment, using a PM30 membrane (Millipore). In the filtration, the solution was concentrated and distilled water was added so that it was possible to reduce the conductivity of the solution to the level required by ion exchange chromatography. The conductivity was measured by using an EDV Instruments conductivity instrument
10 (platinum electrode Mettler Toledo).

After ultrafiltration, the solution was purified by anion exchange chromatography (DEAE Sepharose Fast Flow, h=10 cm, V=20 ml, Pharmacia). The resin was equilibrated at room temperature with 0.01 M acetate buffer, pH 5. The proteins were eluted with an increasing
15 linear saline gradient by using 0-0.5 M sodium sulphate (Merck). The total volume of the gradient was 90 ml and the flow rate 2 ml/min. During the gradient, 4-ml fractions were collected. The protein content, laccase activity, and the conductivity of the fractions were assayed. The laccase activity and the protein content of the fractions are shown in Fig. 2.

20 The best laccase fractions of the anion exchange were combined, and they were further purified by hydrophobic interaction chromatography (HIC) (Phenyl Sepharose Fast Flow, h=9 cm, V=18 ml, Pharmacia). The hydrophobicity of the proteins was increased before the run by adding Na₂SO₄ into the sample, so that the salt content became 0.7 M. The resin was equilibrated with 1 M Na₂SO₄ in 0.02 M citrate buffer, pH 5, at room temperature. The
25 sample was eluted from the column with a decreasing linear saline gradient by using 0.7-0 M Na₂SO₄ in citrate buffer. The total volume of the gradient was 90 ml and the flow rate 2 ml/min. After the gradient, the resin was washed with the equilibrium buffer and, finally, with water. During the gradient and subsequent washes both with buffer and water, 3.5-ml fractions were collected, the laccase activity, the protein content, and the salt content of
30 which were measured (Fig. 3).

The most interesting fractions of both the ion exchange and HIC were analysed by SDS polyacrylamide gel electrophoresis (SDS-PAGE) for monitoring the purification of the laccase in accordance with the method of Laemmli (1970). In the gel electrophoresis, the equipment of Bio-Rad (Bio-Rad Ready Gel Cell), and ready polyacrylamide gels (12 %
5 Tris-HCl Ready Gel) were used. The gels were stained with a Coomassie Brilliant Blue R 350 dye solution (Pharmacia). The Prestained Protein Marker Broad Range #7708S (New England BioLabs) was used as the protein standard.

10 The molecular weight of the *M. albomyces* laccase as defined by the SDS PAGE was 80 kDa.

Example 3.

Characterization of the *M. albomyces* laccase

15

Temperature dependence of the activity

The dependence of activity of the purified *M. albomyces* laccase on the temperature was defined by measuring the laccase activity by ABTS at temperatures of 25, 40, 50, 60, 70, 80 and 90 °C. The enzyme was diluted into tempered 0.025 M succinate buffer, pH 4.5.
20 Immediately after adding the enzyme, the tempered ABTS solution was added. The sample was incubated for two minutes at the desired temperature, after which the absorbance was measured at the wavelength of 436 nm. In order for the added enzyme solution not to considerably decrease the temperature of the buffer, the volume of the enzyme added into the buffer in all dilutions was below 7 % of the total volume. The dependence of the activity
25 of the *M. albomyces* laccase on the temperature is shown in Fig. 4.

pH optimum

The pH optimum of the purified *M. albomyces* laccase was determined by measuring the laccase activity by the guaiacol method in McIlvaine buffer at pH values of 3, 4, 5, 6, 7 and
30 8. Guaiacol was selected because ABTS does not work when the pH is over 7. The enzyme was diluted into the buffers, and immediately after adding the enzyme, 25 µl of a 1% guai-

acol solution was added. The reaction was followed for 5 minutes by a spectrophotometer at the wavelength of 465 nm. The dependence of the activity of the *M. albomyces* laccase on the pH is shown in Fig. 5.

5 Thermal stability

The thermal stability of the *M. albomyces* laccase was determined by incubating the enzyme at 50, 60, 70 and 80°C in 0.06 M citrate buffer, pH 6. The residual activity of the enzyme at each temperature was defined by the ABTS method after incubation of 15, 30, 60 and 120 minutes. The results of the thermal stability measurements are shown in Fig. 6.

10

The thermal stability of the purified *M. albomyces* laccase was determined in the conditions described above. The results are shown in Fig. 7. The results indicate that the residual activity of the enzyme remained essentially unchanged at 60°C for as much as 2 hours. After 4 hours of incubation, there was still 60% residual activity left and after 6 hours, 40%.

15

pH stability

The pH stability of the *M. albomyces* laccase was determined by incubating the enzyme at pH values 2, 3, 4, 5, 6, 7, and 8 in McIlvaine buffer at room temperature. The residual activity of the enzyme in the incubated samples at each pH was defined by the ABTS method after incubation of 1, 3, 5, and 22 hours. The results of the pH stability measurements are shown in Fig. 8.

20

Isoelectric point

The isoelectric point of the *M. albomyces* laccase was determined by isoelectric focusing. The 0.5 mm thick polyacrylamide gel contained 7.5% acrylamide (Merck), 0.225% N,N'-bis-methyleneacrylamide (Merck), 6 % ampholyte (Pharmalyte 2.5 – 5 for IEF, Pharmacia), 0.05% ammonium persulphate (Merck) and 0.05% N,N,N,N'-tetramethylene diamine (Merck). Isoelectric focusing was carried out using the Multiphor II electrophoresis equipment (Pharmacia LKB). The gels were stained using active dyeing, wherein the gel was dipped for about 10 seconds in a dilute ABTS solution (1 g/l). After about 15 minutes, the

25

30

laccase bands appeared green. The pH distribution of the gel was determined using a pH surface electrode (Mettler Toledo). On the basis of isoelectric focusing, the isoelectric point of the *M. albomyces* laccase was 4.0.

5 **Example 4**

Use of the *M. albomyces* laccase in oxidation of dyes

The tests were conducted by using a thermophilic *M. albomyces* laccase and the reference tests by a well-known laccase, which was isolated from *Trametes hirsuta* fungus. A conventional substrate, 2,6-dimethoxyphenol, was used as the substrate. The substrate concentration was 1 mmol/l, pH 4.5-7.5, and the temperatures 40 or 60°C. The amount of laccase was 15 nkat/g. The reaction mixtures were aerated during the test. At the end of the test (30 min), it was discovered that the thermophilic *M. albomyces* laccase worked better at a high temperature and pH, when the formation of colour from the dimethoxyphenol was measured at a wavelength of 468 nm (Fig. 9 and 10).

Laccases can be used in removing colour from textile dyes. Laccases oxidize many dyes, which can be observed as colour removal. The colour removal of textile dyes was measured by using a dye commonly used in the textile industry, Diamond Black PLC. The laccase treatment was carried out at pH 7.5 and temperatures of 40 and 60°C for 2 hours, the laccase dose being 20 nkat/mg of dye. On the basis of a maximum absorbance (560 nm) measurement, the *Melanocarpus* laccase removed 34% of the colour at 40°C and 16% at 60°C, while the *Trametes* laccase only removed 14% at 40°C and only 1% at 60°C (Fig. 1).

The ability of the *M. albomyces* laccase to remove colour was compared with that of the *T. hirsuta* laccase at pH 5 and 8. The colour removal ability was studied using dyes commonly used in the textile industry. A colour solution (50 mg/l) was oxidized with a laccase dose of 1 nkat/ml in 50 mM succinate buffer (pH 5) or in a phosphate buffer (pH 8) for 24 hours at a temperature of 40°C. The colour removal was followed visually. The results are shown in Table 1. As the results indicate, both enzymes oxidized the tested dyes so that they turned colourless at pH 5. The *T. hirsuta* laccase was ineffective at pH 8, and did not

oxidize the dyes tested, unlike the *M. albomyces* laccase, which even at pH 8 was able to oxidize the dyes. The results indicate that the behaviour of the *M. albomyces* laccase in oxidation of dyes is considerably more extensive than that of the conventional *Trametes* laccase.

5

Table 1. Removal of textile dyes with the *T. hirsuta* and the *M. albomyces* laccases at pH 5 and 8. (+=colour removed from the solution, -=no colour removal from the solution)

Dye	<i>T. hirsuta</i> laccase		<i>M. albomyces</i> laccase	
	pH 5	PH 8	pH 5	pH 8
Acid + metal complex dyes				
Acid Blue 113	+	-	+	+
Lanaset Blau 5G	+	-	+	+
After chrome dyes				
Diamond Fast Brown	+	-	+	+
Diamant Schwarz PLC	+	-	+	+

10 Example 5

Use of the M. albomyces laccase in polymerisation

The test studied the polymerisation of a model compound (lignin, Westvaco or Indulin AT, Sigma) with a thermophilic laccase. Lignin is a typical aromatic substrate of laccase. The reference tests were carried out using a laccase that had been isolated from *Trametes hirsuta*. The test conditions were as follows: substrate concentration 1%, pH 7.5, and temperatures 40 or 60°C. The amount of laccase was 200 nkat/g. The reaction mixtures were aerated during the test. The polymerisation was followed by determining the increase in the molecular weight of the polymerised lignin by GPC (gel permeation chromatography). At the end of the reaction (30 minutes), it was discovered that the thermophilic *M. albomyces*

20

laccase worked more effectively than the reference laccase at a high temperature and pH. Fig. 12 illustrates a situation, wherein the polymerisation of lignin is carried out at 40°C and, correspondingly, at 60°C in Fig. 13. The *Melanocarpus* laccase causes an average increase in the molecular size and a decrease in the portion of small molecules obviously better than the *Trametes* laccase.

Example 6

Use of the *M. albomyces* laccase in improving the runnability of paper machines

Laccase can be used to improve the runnability of paper machines by polymerising compounds originating in lignin and extractives and by decreasing the microbiological problems of the paper machine. Generally, the conditions of the paper machine are: the pH is 5-7 and the temperature 60-80°C. Treatment tests were conducted using a thermophilic and a conventional reference laccase in the paper machine conditions by using soluble compounds originating from lignin and extractives, isolated from the paper machine. The substrate concentration was 0.5% dry weight, the pH 7, and the temperature 70°C. Intensive agitation ensured sufficient oxygen content in the reaction. The reaction was monitored as the polymerisation of the aromatic and other compounds, which worked as the substrates of the laccase, by gel filtration (GPC). It was discovered that the reaction by the thermophilic *M. albomyces* laccase was more effective than by the reference laccase.

Fig. 14 shows how the *Melanocarpus* and *Trametes* laccases behave in the polymerisation of compounds originating from lignin and extractives at 70°C. In Fig. 14, the *Melanocarpus* laccase has clearly reduced the amount of small molecules.

Example 7

Use of the *M. albomyces* laccase in oxidation of fibres

Lignin-containing fibres, which were mechanically ground without chemical delignification, were treated with thermophilic *M. albomyces* laccase to activate and polymerise the lignin on the surface of the fibres. The treatments were conducted in the following condi-

tions: the temperatures were 50 and 70°C, the pH 6, and the laccase concentration 200 nkat/g of fibres. The reaction mixture was aerated by blowing oxygen into the reaction mixture. The reference was a conventional laccase that was isolated from the *Trametes hirsuta* fungus. Hand sheets were prepared from the treated fibres. The physical properties of the sheets were measured. The results (Table 2) indicate that the density of the sheets made of the fibres treated with the *M. albomyces* laccase was clearly higher and the light scattering smaller than those of the reference sheets. No corresponding change was observed in sheets, which were made of fibres treated with the *T. hirsuta* laccase. The effect was especially obvious at an increased temperature of 70°C, wherein the *M. albomyces* laccase, because of its good thermal stability, behaved advantageously compared with the reference laccase that was isolated from the *T. hirsuta* fungus. On the basis of the result, it can be concluded that it was possible to use the *M. albomyces* laccase to polymerise lignin onto the surface of the fibres in the conditions used.

Table 2. The density and light scattering of the sheets made from fibres treated with the *M. albomyces* and *T. hirsuta* laccase

Test	T (°C)	Density (kg/m ³)	Light scattering (m ² /kg)
No enzyme	70	339	55.0
<i>T. hirsuta</i> laccase	70	347	55.2
<i>M. albomyces</i> laccase	70	367	53.6

Example 8

Use of the M. albomyces laccase in delignification of fibres

Mediators can be used for expanding the substrate range of the laccase, among others, in delignification or other indirect oxidation of polymers. Hydroxybenzotriazole was selected as the mediator. Thermophilic laccase was used to degrade the lignin of kraft fibres in mediator-assisted oxidation. The reaction conditions were as follows: Kraft pulp, the kappa number about 29, the consistency 4%, the amount of mediator 1% of the pulp, the pH ad-

justed to 7, and the temperature 70°C, the reaction time 2 hours, the amount of laccase 500 nkat/g, in oxygen pressure (0.5 mPa). *Trametes* laccase served as a reference material. After the treatments, the pulps were subjected to alkali extraction, chelation, and one-stage peroxide bleaching. Dissolution of the lignin was monitored measuring the absorbance of the filtrate at a wavelength of 280 nm after alkali extraction. Handsheets were prepared of the pulps and ISO Brightness, and the kappa numbers were measured. As the results indicate (Table 3), the *M. albomyces* laccase works more effectively in the test conditions than the reference laccase; it decreased the kappa value of the sheets (the amount of lignin) and increased the absorbance_{280nm} of the filtrate more than the reference laccase.

Table 3. The effect of LM treatment on the bleachability of kraft pulp (κ 29) with an LQ-E-P sequence

Test	A280 from filtrate	Kappa value
No enzyme	11.9	18.9
<i>T. hirsuta</i> laccase	14.3	16.3
<i>M. albomyces</i> laccase	32.7	15.4

Example 9

Isolation of the gene and the cDNA of the *M. albomyces* laccase

Total DNA was isolated from the cells according to Raeder and Broda (1985). A genomic library was constructed using a commercial reaction package (SuperCos I Cosmid Vector Kit, Stratagene), following the instructions of the manufacturer. 100 μ g of DNA were partially cleaved with 5 U of a *Sau*3AI restriction enzyme (New England Biolabs) by incubating for 10 minutes at 37°C. The digested DNA was dephosphorylated with 20 U of CIAP (Calf Intestinal Alkaline Phosphatase, Finnzymes). The DNA molecules of different lengths were separated with a 15-30% sucrose gradient by ultracentrifugating for 22 hours at a temperature of 20°C and a rotational speed of 22 000 rpm (the Beckman SW 41 TI ro-

tor). The 12-ml sucrose gradient was divided into 300- μ l fractions, and 10 μ l of every second fraction were examined on 0.5% agarose gel by electrophoresis. The fractions, which on the basis of the gel electrophoresis contained DNA fragments of more than 20 kilobase-pairs in length, were combined, and the DNA was precipitated from them with ethanol. The
5 obtained DNA (about 2 μ g) was inserted into SuperCos I cosmid vector (about 1 μ g), which had been digested with XbaI (New England Biolabs), and thereafter dephosphorylated with CIAP and, finally, digested with BamHI (Boehringer Mannheim). Ligation was carried out with T4 DNA ligase (Promega) by incubating the mixture at 4°C overnight. The ligation mixture was packed into λ particles by using a commercial reaction package
10 (Gigapack III Gold packaging extract, Stratagene), following the instructions of the manufacturer, and the packed phage was used to infect *E. coli* host cells (the XL1-Blue MR strain, Stratagene). One ligation provided more than 5×10^6 clones, which can be considered a representative gene bank.

15 The hybridisation probe for the screening of the gene bank was obtained from the *lac2* gene of the *Podospora anserina* fungus. This gene was selected as the probe, because the N-terminal amino acid sequence and the internal peptides of the *M. albomyces* laccase were homologous to the amino acid sequence of the *P. anserina* laccase. *P. anserina* fungus was grown on a substrate, which is described in Fig. 1. Genomic DNA was isolated from a mycelium, which was collected and freeze-dried after three days of growing, using a commercial reaction package (Easy DNA Kit, Invitrogen) in accordance with the manufacturer's
20 instructions. The *lac2* gene was multiplied by a PCR reaction, using primers that are based on the published sequence of the *lac2* gene: 5'-TGCCACACTGCCGCCAACCGTGCT-3' (SEQ ID NO: 3) (forward) and 5'-GTTCTTGATATACCAATCAGGATG-3' (SEQ ID
25 NO: 4) (reverse). The PCR program used for multiplication comprised 26 cycles, wherein the temperature program was as follows: denaturation of DNA at 94°C for 45 seconds, insertion of primers at 55°C for 1 min, extending the DNA chain with polymerase at 72°C for 2.5 min. Finally, the chain was extended at 72°C for 5 minutes. The obtained fragment of about 1.9 kilobasepairs in length was purified by agarose gel electrophoresis. Its behaviour as a probe was examined by Southern hybridisation with the genomic DNA of the *M.*
30

albomyces fungus as follows: in two different reactions, 40 µg of DNA was cleaved with 80 U of EcoRI and HindIII restriction enzymes (New England Biolabs) at 37°C for 5 hours. The fragments were separated by electrophoresis on 0.8% agarose gel; DNA was denatured and transferred onto Hybond N membranes (Amersham Pharmacia Biotech) (the method is described in Sambrook *et al.*, 1989). The *P. anserina lac2* gene was labelled with α -³²P-dCTP by using a commercial reaction package (Random primed DNA labelling kit, Boehringer Mannheim) in accordance with the manufacturer's instructions, and the bonding of the probe was examined at four different hybridisation temperatures: 48, 50, 55, and 60°C. The hybridisation solution contained 6xSSC, 1x Denhardt's (Sambrook *et al.*, 1989), 0.5% SDS, and 100 µg/ml of Herring Sperm DNA (the SSC contains 0.15 M NaCl and 0.015 M sodium citrate at pH 7.0). The hybridisation contained about 5x10⁵ cpm/ml of labelled probe. After hybridisation, the membranes were washed in 2xSSC containing 0.1% SDS twice for five minutes at room temperature and, after this, for 30 minutes at the same temperature as the hybridisation was made at. The membranes were enclosed in an exposure cassette with a film, and the exposed film showed that the *P. anserina lac2* gene was hybridised with a DNA fragment, which was about 4.5 kilobasepairs in length and cleaved with ExoRI enzyme.

About 5x10⁵ clones from the obtained genomic gene bank were placed on agar plates, and colonies that had been growing overnight were transferred to nitrocellulose membranes (Protran, Schleicher & Schuell). The DNA contained by the colonies was denatured (Sambrook *et al.*, 1989) and then attached to the membranes by heating at 80°C for 2 hours. After the attachment of DNA, the residuals of the bacterial colonies were washed from the membranes by scrubbing them with the washing fluid (Sambrook *et al.*, 1989) at 48°C. The DNA bound to the membranes was hybridised overnight at a temperature of 57°C with a *P. anserina lac2* gene that was labelled with α -³²P-dCTP (a hybridisation solution, as above). On the basis of the obtained hybridisation signals, several colonies were picked from the original plates, to which colonies the *P. anserina lac2* gene had been hybridised. These colonies were further hybridised with the *P. anserina lac2* gene and, on the basis of radioactive signals, 6 colonies were selected for further examination. Cosmids (Plasmid purifi-

cation protocol, Tip-500, QIAGEN) were isolated from them. The restriction enzyme, which made it possible to isolate the laccase gene from the cosmid, was sought for by cleaving the cosmids with 19 different restriction enzymes. The obtained fragments were subjected to Southern hybridisation at a temperature of 57°C, using the *P. anserina lac2* gene, as above. An EcoRI fragment of about 4.5 kilobasepairs in length was again hybridised with the *lac2* gene. The cosmid was cleaved with EcoRI, and the fragment was purified by agarose gel electrophoresis. The obtained fragment was inserted into plasmid pBluescriptSK⁻ (Stratagene), and the plasmid was transformed into *E. coli* host cells (strain DH5 α , Gibco BRL) by electroporation. A clone was obtained from the transformation, containing the desired EcoRI fragment in pLLK1 plasmid. A laccase gene was sequenced from this plasmid, using synthesized oligonucleotide primers. The sequencing reactions were carried out using a commercial reaction package (DNA Sequencing Kit, dRhodamine Terminator Cycle Sequencing Ready Reaction, PE Biosystems) in accordance with the manufacturer's instructions.

To establish the introns contained by the *M. albomyces* laccase gene, a complementary DNA (cDNA) corresponding to the *M. albomyces* laccase was cloned by RACE-PCR, using a commercial reaction package (FirstChoiceTM RLM-RACE Kit, Ambion, Inc.) in accordance with the manufacturer's instructions. RNA was isolated from the *M. albomyces* cells by a commercial reaction package (TRIZOL[®] Reagent, Life Technologies) in accordance with the manufacturer's instructions. The RNA was dephosphorylated and transcribed into DNA by using an inverse transcribing enzyme, whereafter the 5' and 3' ends of the gene were multiplied by separate PCR reactions in accordance with the manufacturer's instructions. The multiplication of both cDNA ends was carried out by two sequential PCR reactions using different primers. In the first reaction, the desired part of the laccase cDNA was multiplied using a gene-specific primer, and the second reaction ensured the multiplication of the laccase cDNA by using another gene-specific primer. In each PCR reaction, one of the primers needed for the multiplication of the DNA came from the RLM-RACE Kit reaction package, and it was bound to the adapter regions connected to the ends of the cDNA. The PCR program used for multiplication comprised 35 cycles, wherein the temperature program was as follows: denaturation of DNA at 94°C for 30 seconds, insertion of

primers at 60-62°C for 30 sec, extension of DNA chain at 72°C for 2 minutes. In the multiplication of the 5' end, the insertion temperature of the primers was 60°C, and in that of the 3' end, 62°C. Finally, chain extension was carried out at 72°C for 7 minutes. The gene-specific primers used in the multiplication of the 5' end were as follows: in the first PCR reaction, 5'-GCCGGTGAGGATGTAGTCGATGAT-3' (SEQ ID NO: 5), and in the second reaction, 5'-AGGTGACGTTGAACCAAGTAGTTGTC-3' (SEQ ID NO: 6). The gene-specific primers used in the multiplication of the 3' end were as follows: in the first PCR reaction, 5'-CTGGTGCACTTCACGCAGAACAA-3' (SEQ ID NO: 7), and in the second reaction, 5'-AGAACCACTTCCAGGTGTCGCT-3' (SEQ ID NO: 8). From the RLM-RACE reactions, a fragment of 1194 basepairs in length was obtained from the 5' end, and from the 3' end, a fragment of 1322 basepairs in length. The fragments were isolated by agarose gel electrophoresis and cloned into a pCR2.1-TOPO™ vector by using a commercial reaction package (TOPO TA Cloning Kit, Invitrogen) in accordance with the manufacturer's instructions. The plasmid was transformed into *E. coli* host cells (strain TOP10F', Invitrogen) by using electroporation in accordance with the instructions of the reaction package. The cloned fragments were sequenced and the positions of the introns in the gene of the *M. albomyces* laccase were established by comparing the genomic sequence with the cDNA sequence obtained.

The genomic sequence of the gene is shown in Fig. 15. The introns are underlined.

References:

Altschul et al. 1990. J. Mol. Biol. 215, 403-410.

Berka et al. 1997. Applied and Environmental Microbiology 63(8):3151-3157.

Bharathi, M.S. and Ramalingam, K. 1993. J. Anim. Morphol. Physiol. 40(1&2):153.

Diamantidis et al. 2000. Soil Biology & Biochemistry 32: 919-927.

Fernandez-Larrea & Stahl. 1996. Mol. Gen. Genet. 252, 539-551.

Heinzkill et al. 1998. Appl. Environ. Microbiol. 64: 1601 – 1606.

Jönsson et al. 1997. Curr. Genet. 32: 425 – 430.

Laemmli 1970. Nature 227: 680-685.

Lowry et al. 1951. *J. Biol. Chem.* 193:265-275.

Kojima *et al.* 1990. *J. Biol. Chem.* 265: 15224 – 15230.

Niku-Paavola et al. 1988. *Biochem. J.* 254:877-884

Paszczynski et al. 1985. *FEMS Microbiol. Lett.* 29 (1985) 37 – 41. Ravanko, K. 1996.

- 5 Termostabiilin lakkaasin seulonta (Screening of Thermostable Laccase). Master's Thesis.
97 p. 4 Appendices. (In Finnish)

Raeder and Broda, 1985. *Lett. Appl. Microbiol.* 1: 17-20.

Saloheimo *et al.* 1985. *J. Gen. Microbiol.* 137: 1537-1544.

Saloheimo and Niku-Paavola. 1991. *Bio/Technology* 9: 987 – 990.

- 10 Sambrook *et al.*, 1989, *Molecular cloning: a laboratory manual*, 2nd. edn. Cold Spring
Harbor Laboratory Press, Cold Spring Harbor, New York.
Yaver *et al.* 1996., *Appl. Environ. Microbiol.* 62: 834 – 841.

Claims

1. A laccase enzyme, characterized in that it is isolatable from the strains of the *Melanocarpus* genus, and that the pH optimum of the enzyme is within 5-8 and the enzyme
5 functions at pH 3-9 and a temperature of 30-80°C.

2. The laccase enzyme according to claim 1, characterized in that it is isolatable from the *M. albomyces* strain.

10 3. The laccase enzyme according to claim 1 or 2, characterized in that the pH optimum of the enzyme is within 6-8.

4. The laccase enzyme according to any one of claims 1 to 3, characterized in that the enzyme works best at a temperature of 50-80°C.

15 5. The laccase enzyme according to any one of claims 1 to 4, characterized in that the pH optimum of the enzyme is about 7.5 and the temperature optimum about 70°C.

20 6. The laccase enzyme according to any one of claims 1 to 5, characterized in that the activity of the enzyme remains essentially unchanged for 2 hours at 60°C.

7. The laccase enzyme according to any one of claims 1 to 6, characterized in that the isoelectric point of the purified enzyme is about 4.0 as determined by isoelectric focusing, and its molecular weight is about 80 kDa as determined by SDS-PAGE.

25 8. An isolated and purified *Melanocarpus* laccase enzyme, preferably a *Melanocarpus albomyces* laccase enzyme.

30 9. An enzyme preparation comprising *Melanocarpus* laccase, preferably *Melanocarpus albomyces* laccase enzyme as the main activity.

10. An enzyme preparation, characterized in that it comprises more laccase according to any of the preceding claims than what the strains of the *Melanocarpus* genus naturally produce into their culture solution.

5 11. The preparation according to claim 9 or 10, characterized in that the pH optimum of the enzyme is within 5-8 and that the enzyme works at pH 3-9 and a temperature of 30-80°C.

10 12. The preparation according to any one of claims 9 to 11, characterized in that the isoelectric point of the enzyme is about 4.0 as determined by isoelectric focusing and that its molecular weight is about 80 kDa as determined by SDS-PAGE.

13. A nucleic acid molecule that encodes a polypeptide that has laccase enzyme activity, characterized in that the nucleic acid molecule is selected from the group comprising:
15

a) a nucleic acid molecule comprising the coding region of the nucleotide sequence as depicted in SEQ ID NO:1 or Fig. 15;

b) a nucleic acid molecule that encodes a polypeptide comprising the amino acid sequence as depicted in SEQ ID NO:2 or Fig. 15;

20 c) a nucleic acid molecule comprising a coding sequence that differs from the coding sequence of SEQ ID NO:1 or Fig. 15 due to the degeneracy of the genetic code;

d) a nucleic acid molecule that hybridizes to the nucleotide sequence of SEQ ID NO:1 or Fig. 15; and

e) a nucleic acid molecule that encodes a polypeptide, which has laccase activity
25 and the amino acid sequence of which shows at least 73% identity with the amino acid sequence of SEQ ID NO:2 or Fig. 15.

14. The nucleic acid molecule that encodes a polypeptide that has laccase enzyme activity, characterized in that the amino acid sequence of the polypeptide shows at least 80%
30 identity with the amino acid sequence of SEQ ID NO:2 or Fig. 15.

15. The nucleic acid molecule according to claim 13 or 14, characterized in that it is a nucleic acid molecule which encodes a polypeptide that comprises the amino acid sequence as depicted in SEQ ID NO:2 or Fig. 15.

5 16. The nucleic acid molecule according to any one of claims 13 to 15, characterized in that it is a nucleic acid molecule that comprises the coding region of the nucleotide sequence as depicted in SEQ ID NO:1 or Fig. 15.

10 17. The nucleic acid molecule according to any one of claims 13 to 16, characterized in that it encodes the enzyme according to any one of claims 1 to 8.

18. A vector, characterized in that it comprises the nucleic acid molecule according to any one of claims 13 to 17.

15 19. A micro-organism host, characterized in that the nucleic acid molecule according to any one of claims 13 to 17 or the vector according to claim 18 has been transferred into it.

20 20. The host according to claim 19, characterized in that it is a filamentous fungus, yeast or bacterium.

21. The host according to claim 19 or 20, characterized in that it belongs to the genera *Aspergillus*, *Trichoderma* or *Fusarium*.

25 22. A method for producing laccase, characterized in

- transferring a nucleic acid molecule according to any one of claims 13 to 17 or the vector according to claim 18 into a micro-organism host cell for expressing the
- nucleic acid molecule and, optionally, for secreting it from the host cell; and
- recovering the polypeptide having laccase activity either from the cells or from the

30 culture solution of the micro-organism host.

23. A polypeptide, which has laccase activity, c h a r a c t e r i z e d in that it is encoded by a nucleic acid molecule, which is selected from the group comprising:

- a) a nucleic acid molecule which comprises the coding region of the
- b) nucleotide sequence as depicted in SEQ ID NO:1 or Fig. 15;
- 5 c) a nucleic acid molecule that encodes a polypeptide which comprises the amino acid sequence as depicted in SEQ ID NO:2 or Fig. 15;
- d) a nucleic acid molecule that differs from the coding sequence of the
- e) nucleotide sequence of SEQ ID NO:1 or Fig. 15 due to the degeneracy of the genetic code;
- 10 f) a nucleic acid molecule which hybridizes to the nucleotide sequence of SEQ ID NO:1 Fig. 15; and
- g) a nucleic acid molecule that encodes a polypeptide which has laccase activity and the amino acid sequence of which shows at least 73% identity with the amino acid sequence of SEQ ID NO:2 or Fig. 15.

15

24. A polypeptide having laccase activity, c h a r a c t e r i z e d in that it comprises an amino acid sequence which shows at least 80% identity with the amino acid sequence of SEQ ID NO:2 or Fig. 15.

20

25. An enzyme preparation, c h a r a c t e r i z e d in that it comprises, as the main activity, the polypeptide according to claim 23 or 24, which has laccase activity.

26. An enzyme preparation, c h a r a c t e r i z e d in that it comprises more of the polypeptide according to claim 22 or 23 which has laccase activity, than what the strains of the
25 *Melanocarpus* genus naturally produce into their culture solutions.

27. The enzyme preparation according to any one of claims 9 to 12, 25 or 26, c h a r a c -
t e r i z e d in that it comprises more than 10mg/l, preferably ≥ 30 mg/l of laccase.

30

28. The enzyme preparation according to any one of claims 9 to 12, 25 to 27, c h a r a c -
t e r i z e d in that it comprises at least one additive, such as a stabilizer and/or a buffer.

29. The use of the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 in oxidizing colouring agents.

30. The use of the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 in polymerising compounds.

31. The use of the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 in oxidizing fibres.

32. The use of the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 in delignification of fibres.

33. The use of the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 in improving the runnability of paper machines.

34. A method for oxidizing colouring agents, characterized in that, in the presence of a sufficient amount of oxygen, the colouring agent is brought into contact with the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 at pH 4-8, a temperature of 40-80°C, for 5

min – 24 hours with an amount of laccase, which is about 1 - 1000 nkat/g of colouring agent.

35. A method for polymerising compounds, characterized in that, in the presence of
5 a sufficient amount of oxygen, the compound to be polymerised is brought into contact with the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 at pH 4-8, a temperature of 40-80°C, for 5 min – 24 hours with an amount of laccase, which is about 1-1000 nkat/g of the
10 substance that is to be polymerised.

36. A method for oxidizing fibres, characterized in that, in the presence of a sufficient amount of oxygen, the fibres that are to be oxidized are brought into contact with the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one
15 of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 at pH 5-8, a temperature of 50-80°C, for 5 min – 24 hours with an amount of laccase, which is about 1 - 1000 nkat/g of dry matter fibres.

20 37. A method for delignifying fibres, characterized in that, in the presence of a sufficient amount of oxygen, the fibre that is to be delignified is brought into contact with the laccase according to any one of claims 1 to 8, the enzyme preparation according to any one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22, the pH being within 5-8, the temperature
25 within 50-80°C, for 5 min – 24 hours with an amount of laccase, which is about 1 - 1000 nkat/g of dry matter fibres.

38. A method for improving the runnability of paper machines, characterized in that the laccase according to any one of claims 1 to 8, the enzyme preparation according to any
30 one of claims 9 to 12 or 25 to 28, or the polypeptide according to claim 23 or 24, or the laccase obtained by the method according to claim 22 is added into the process water or the

head box of the paper machine 1-1000 nkat/g of dry matter, fibres or litres of circulation water.

39. A method according to any one of claims 34 to 38, c h a r a c t e r i z e d in that a me-
5 diator is used in the method, when needed.


1/10

Fig. 1




Fig. 2

2/10




Fig. 3




Fig. 4

3/10




Fig. 5




Fig. 6

4/10




Fig. 7


5/10

Fig. 8

6/10




Fig. 9




Fig. 10

7/10




Fig. 11




Fig. 12

8/10




Fig. 13




Fig. 14

9/10

atgaagaccttcaccagcgccctggcgctcgtgggtgggcatgcttgccccgggtgccgtc
M K T F T S A L A L V V G M L A P G A V
gttgccgcgcctcccagcactccggcccagcgagatctggctcgagctgcgcgagggcagg
V A A P P S T P A Q R D L V E L R E A R
caggagggcggaaggacctccgtccccgtgagccgacgtgcaacacgcccagcaaccgg
Q E G G K D L R P R E P T C N T P S N R
gcgtgctggagcgacggcttcgacatcaacaccgactacgaagtcagcacgcccggatacc
A C W S D G F D I N T D Y E V S T P D T
ggcgtcactcagtctgtgagtcctcctcctccgc
G V T Q S

ccccttggtttcgggaaggctgtccgaagggaaggaaagtccgaggctaactgcagacag

tacgtattcaacctcactgaggttgacaactggatgggcccagacggcgtcgtcaaggag
Y V F N L T E V D N W M G P D G V V K E
aaggatgattgatcaatgggtgagtcgaccgcgaatgggatgtcgagttgccatccaaaa
K V M L I N

tagacctgtgtaactgactgggttcgatcacagggaacattatgggtacgtcttatctg

G N I M

gctgtcccggcacgccaatggcctttcttttcccttgagcgtaggacgtgcgacggactga

ctgagaacaataggccccaacatcgctcggaactggggtgatacggctcgaggtcacctgtg

G P N I V A N W G D T V E V T V

atcaacaaccttgtgaccaacgga acgtcgatccactggcacggcatccaccagaaggac
I N N L V T N G T S I H W H G I H Q K D
accaacctgcacgacggcgccaacggcggtgaccgagtgctccgatccccgccaaggcgggc
T N L H D G A N G V T E C P I P P K G G
cagcggacgtaccgtggcgggcgcgccagctggcaccagctggtagcactcgcacttc
Q R T Y R W R A R Q Y G T S W Y H S H F
tcggcgacgtacggcaacggcggtgggtgggcacgatccagatcaacggccccggcgctcgtg
S A Q Y G N G V V G T I Q I N G P A S L
ccctacgacatcgaccttggcggtgtccccatcaccgactactactaccgggcccggccgac
P Y D I D L G V F P I T D Y Y Y R A A D
gacctgggtgcacttcacgcagaacaacgcgcgcccttcagcgacaacgtgctcatcaac
D L V H F T Q N N A P P F S D N V L I N
ggcacggccgtcaacccgaacacggggcgagggccagtagcgaacgtgacgctgacgccg
G T A V N P N T G E G Q Y A N V T L T P
ggcaaggcgacccgcctcgcatcctcaacacgtcgaccgagaaccacttccaggtgtcg
G K R H R L R I L N T S T E N H F Q V S
ctcgtcaaccacaccatgacgggtcatcgccgcgacatgggtgcccgtcaacgccatgacg
L V N H T M T V I A A D M V P V N A M T
gtcgacagcctgttcctggccgtcgggcagcgctacgacgtcgtcatcgacgcctcgaga
V D S L F L A V G Q R Y D V V I D A S R
gccccggacaactactggttcaacgtcacctttggcgggccaggcggtgcgggcggtcg
A P D N Y W F N V T F G G Q A A C G G S
ctcaaccgcgacccggccgcatcttccactacgcggcgcccgggcggtgcccacc
L N P H P A A I F H Y A G A P G G L P T
gacgagggcacgccccgggtcgaccaccagtgccctggacacgctcgacgtgcgccccgtc
D E G T P P V D H Q C L D T L D V R P V
gtgcccgcgacgctgcccgtcaacagcttcgtcaagcggccccgacaacacgctgcccgggtg
V P R S V P V N S F V K R P D N T L P V
gcgctcgacctgaccggcacgccccgtgttcgtgtgg

A L D L T G T P L F V W

aaggtcaacggcagcgacatcaacgtcgactgggggaagcccatcatcgactacatcctc
K V N G S D I N V D W G K P I I D Y I L
accggcaacaccagctacccccgtgtcggaacaacatcgatgcaggttgatgcgctcgatcag
T G N T S Y P V S D N I V Q V D A V D Q

10/10

gtatgtcctctgttaaagccttacgtcgtacgattgcgctggcaaatacacacctcgtact
gacgccaaaccaccacctcccttctagtggacatactggctcatcgagaacgatccggag
W T Y W L I E N D P E
ggcccttcagcctgccgcacccgatgcacctccac
G P F S L P H P M H L H
gtaagtggccaaccctgtcacgtac
gttgcaaccgcctctctccggggccccccaactcacacttcggactcccgtcccacagggc
G
cacgacttcctcgtgctggggcggtcgcccgacgtgccggcggtcgcagcagcgcttc
H D F L V L G R S P D V P A A S Q Q R F
gtgttcgacccggcgtggacctggcgcggtcaacggcgacaacccgccgcggcgac
V F D P A V D L A R L N G D N P P R R D
accacgatgctgccggcggtgggtgctgctcgccttcgcaccgacaacccgggc
T T M L P A G G W L L L A F R T D N P G
gcctggctcttccactgccacatcgctggcacgtgtcgggcggcctgtcggtcgacttc
A W L F H C H I A W H V S G G L S V D F
ctcgagcgccccgccgacctgcgccagcgcacatctcccaggaggacgaggacgacttcaac
L E R P A D L R Q R I S Q E D E D D F N
cgcgtctgcgacgagtgccgcgcctactggccgacgaatccctacccaagatcgactcg
R V C D E W R A Y W P T N P Y P K I D S
ggcctgaagcgtcgccgctgggtggaggagcgcgagtggtggttcggttgatgggggaaa
G L K R R R W V E E S E W L V R -
gggggggtaggtgcgattcaggggtacctaggggtgcacttgatggtgatgctcgatggag
aattggttttggttacttggtgttcactttcgacatgcgttggtccttggttgatgtttt
taggttggtccagatgatggatgatatggtaccgaggggaactgtggtccttgcttcgaaag
gggacttcattgttatggtaccacaggaccagttatcgaagcatccttggtttcaatcgca poly-A
tcttttttccccgatgcctcgagtacatgatgcccatagtgagtcgtagcacaccagcc begins
acaccacctacctcccttcccgcaacgccatcatgcggtacgccaaatcaaggctctcca
caatcccatgccaatgcgcacctccgcacagcaacctcgatcc

FIG. 15

SEQUENCE LISTING

<120> Novel laccase enzyme and the gene encoding the enzyme

<130> vtt105

<140>

<141>

<150> 20001240

<151> 2000-05-23

<160> 8

<170> PatentIn Ver. 2.1

<210> 1

<211> 3116

<212> DNA

<213> Melanocarpus albomyces

<400> 1

```

gaattcagga tggcatcgag tgaagggcga ggaggatgtg ccgaatacga gctcgtcgac 60
ggcttcctgc tccgtcctgc gctctcctgg gatcgagcgg catggggatt tcttctatat 120
aaggggctcg acagtcgcga tcatgagatt gattttttcca cctcaccag gcacaagcca 180
gccatcagaa cctctcttcc accttcatca ggcaactctcc ttccgtctgt cgagctcttt 240
cccacatttc tagcaggcgg tttegcacacc agccgtcttc acaccatgaa gaccttcacc 300
agcgccctgg cgctcgtggt gggcatgctt gccccgggtg ccgtcgttgc cgcgccctcc 360
agcactccgg cccagcgaga tctggtcgag ctgcgcgagg cgaggcagga gggcggaag 420
gacctccgtc cccgtgagcc gacgtgcaac acgcccagca accgggctg ctggagcgac 480
ggcttcgaca tcaacaccga ctacgaagtc agcacgccgg ataccggcgt cactcagtct 540
gtgagtcctc ctctccgccc ccttggtttc ggggaaggctg tccgaaggga aggaaagtcc 600
gaggctaact gcagacagta cgtattcaac ctactgagg ttgacaactg gatgggcca 660
gacggcgctg tcaaggagaa ggtgatgttg atcaatggtg agtcgaccgc gaatgggatg 720
tcgagttgcc atccaaaata gacctgtgta actgactggt tcgatcacag ggaacattat 780
gggtacgtct tatctggctg tcccggcacg ccaatggcct ttcttttcc tgcgcgtagg 840
acgtgcgacg gactgactga gaacaatagg cccaacatc gtcgcgaact ggggtgatac 900
ggtcgaggtc accgtgatca acaaccttgt gaccaacgga acgtcgatcc actggcacgg 960
catccaccag aaggacacca acctgcacga cggcgccaac ggcgtgaccg agtgtccgat 1020
cccggccaag ggcggccagc ggacgtaccg ctggcgggcg cggcagtatg gcaccagctg 1080
gtaccactcg cacttctcgg cgcagtacgg caacggcgctg gtgggcacga tccagatcaa 1140
cggcccgggc tcgctgccct acgacatcga ccttggcggt ttccccatca ccgactacta 1200
ctaccggggc gccgacgacc tgggtgcact cacgcagaac aacgcgccgc ccttcagcga 1260
caacgtgctc atcaacggca cggccgtcaa cccgaacacg ggcgagggcc agtacgcaa 1320
cgtgacgctg acgcccggca agcggcacgc cctgcgcata ctcaacacgt cgaccgagaa 1380
ccacttccag gtgtcgtcgt tcaaccacac catgacgggtc atcgccgccg acatggtgcc 1440
cgtcaacgcc atgacggtcg acagcctgtt cctggccgct ggccagcgct acgacgtcgt 1500
catcgacgcc tcgagagccc cggacaacta ctggttcaac gtcacctttg gcggccaggc 1560
ggcgtgcggc ggctcgtcga acccgacccc ggccgccatc ttccactacg ccggcgcgcc 1620
cggcggcctg cccaccgacg agggcacgcc cccggtcgac caccagtgcc tggacacgct 1680
cgacgtgcgc cccgtcgtgc cgcgcagcgt gcccgtcaac agcttcgtca agcgggcccga 1740
caacacgctg ccggtggcgc tcgacctgac cggcacgccc ctgttcgtgt ggaaggtcaa 1800
cggcagcgac atcaacgtcg actggggcaa gccatcatc gactacatcc tcaccggcaa 1860
caccagctac cccgtgtcgg acaacatcgt gcaggttgat gccgtcgatc aggtatgtcc 1920
tctgttaaag ccttacgtcg tacgattgcg ctggcaaatc acacctcgta ctgacgcaa 1980
accaccacct cccttctagt ggacatactg gctcatcgag aacgatccgg agggccctt 2040
cagcctgccg caccgatgc acctccacgt aagtggccaa ccctgtcacg tacgttgcaa 2100
ccgcccctct cggggccccc caactcacac ttcggaactc cgtcccacag ggccacgact 2160
tctcgtgct ggggcggctg cccgacgtgc cggcgggcgt gcagcagcgc ttcgtgttcg 2220

```

```

acccggccgt ggacctggcg cggctcaacg gcgacaaccc gccgcgccgc gacaccacga 2280
tgctgccggc cggcggtcgg ctgctgctcg ccttcgcgac cgacaaccgc ggcgccctggc 2340
tcttccactg ccacatcgcc tggcacgtgt cgggcggcct gtcggtcgac ttcctcgagc 2400
gccccgccga cctgcgccag cgcctctccc aggaggacga ggacgacttc aaccgcgtct 2460
gcgacgagtg gcgcgcctac tggccgacga atccctaccc caagatcgac tcgggcctga 2520
agcgtcgccg ctgggtggag gagagcgagt ggctgggttc ttgatggggg aaaggggggg 2580
taggtgcgat tcaggggtac ctagggtgca cttgatgttg atgctcgatg gagaattggg 2640
tttggttact tgttggtcac ttctgacatg cgttggtcct tgtttggatt ttttaggttg 2700
tccagatgat ggatgatatg gtaccgaggg aactgtggtc ttgccttcga aaggggactt 2760
catgttatgg taccacagga ccagttatcg aagcatcctt gttttcaatc gcatcttttt 2820
tcccccgatg cctcgagtac atgatgccca tagtgagtcg tagcacacca gccacaccac 2880
ctacctccct tcccgcacg ccacatcgcg gtacgccaaa tcaaggtcct ccacaatccc 2940
atgccaatcg cgcacacctc gcacagcaac ctcgatcctc ctccggaacg cctcgccagc 3000
atcaatcgcc cgcccaaccc cccgtacatc accaccagc gaaggctccg tctcgtccac 3060
cccaaccacc cgccagtacc gcagcagccc gacgtcctct cgaccggcgc aagcac 3116

```

<210> 2

<211> 623

<212> PRT

<213> *Melanocarpus albomyces*

<400> 2

```

Met Lys Thr Phe Thr Ser Ala Leu Ala Leu Val Val Gly Met Leu Ala
  1                      5                      10                      15

Pro Gly Ala Val Val Ala Ala Pro Pro Ser Thr Pro Ala Gln Arg Asp
                20                      25                      30

Leu Val Glu Leu Arg Glu Ala Arg Gln Glu Gly Gly Lys Asp Leu Arg
    35                      40                      45

Pro Arg Glu Pro Thr Cys Asn Thr Pro Ser Asn Arg Ala Cys Trp Ser
    50                      55                      60

Asp Gly Phe Asp Ile Asn Thr Asp Tyr Glu Val Ser Thr Pro Asp Thr
    65                      70                      75                      80

Gly Val Thr Gln Ser Tyr Val Phe Asn Leu Thr Glu Val Asp Asn Trp
                85                      90                      95

Met Gly Pro Asp Gly Val Val Lys Glu Lys Val Met Leu Ile Asn Gly
    100                      105                      110

Asn Ile Met Gly Pro Asn Ile Val Ala Asn Trp Gly Asp Thr Val Glu
    115                      120                      125

Val Thr Val Ile Asn Asn Leu Val Thr Asn Gly Thr Ser Ile His Trp
    130                      135                      140

His Gly Ile His Gln Lys Asp Thr Asn Leu His Asp Gly Ala Asn Gly
    145                      150                      155                      160

Val Thr Glu Cys Pro Ile Pro Pro Lys Gly Gly Gln Arg Thr Tyr Arg
                165                      170                      175

Trp Arg Ala Arg Gln Tyr Gly Thr Ser Trp Tyr His Ser His Phe Ser
    180                      185                      190

Ala Gln Tyr Gly Asn Gly Val Val Gly Thr Ile Gln Ile Asn Gly Pro
    195                      200                      205

```

Ala	Ser	Leu	Pro	Tyr	Asp	Ile	Asp	Leu	Gly	Val	Phe	Pro	Ile	Thr	Asp	210	215	220
Tyr	Tyr	Tyr	Arg	Ala	Ala	Asp	Asp	Leu	Val	His	Phe	Thr	Gln	Asn	Asn	225	230	235 240
Ala	Pro	Pro	Phe	Ser	Asp	Asn	Val	Leu	Ile	Asn	Gly	Thr	Ala	Val	Asn	245	250	255
Pro	Asn	Thr	Gly	Glu	Gly	Gln	Tyr	Ala	Asn	Val	Thr	Leu	Thr	Pro	Gly	260	265	270
Lys	Arg	His	Arg	Leu	Arg	Ile	Leu	Asn	Thr	Ser	Thr	Glu	Asn	His	Phe	275	280	285
Gln	Val	Ser	Leu	Val	Asn	His	Thr	Met	Thr	Val	Ile	Ala	Ala	Asp	Met	290	295	300
Val	Pro	Val	Asn	Ala	Met	Thr	Val	Asp	Ser	Leu	Phe	Leu	Ala	Val	Gly	305	310	315 320
Gln	Arg	Tyr	Asp	Val	Val	Ile	Asp	Ala	Ser	Arg	Ala	Pro	Asp	Asn	Tyr	325	330	335
Trp	Phe	Asn	Val	Thr	Phe	Gly	Gly	Gln	Ala	Ala	Cys	Gly	Gly	Ser	Leu	340	345	350
Asn	Pro	His	Pro	Ala	Ala	Ile	Phe	His	Tyr	Ala	Gly	Ala	Pro	Gly	Gly	355	360	365
Leu	Pro	Thr	Asp	Glu	Gly	Thr	Pro	Pro	Val	Asp	His	Gln	Cys	Leu	Asp	370	375	380
Thr	Leu	Asp	Val	Arg	Pro	Val	Val	Pro	Arg	Ser	Val	Pro	Val	Asn	Ser	385	390	395 400
Phe	Val	Lys	Arg	Pro	Asp	Asn	Thr	Leu	Pro	Val	Ala	Leu	Asp	Leu	Thr	405	410	415
Gly	Thr	Pro	Leu	Phe	Val	Trp	Lys	Val	Asn	Gly	Ser	Asp	Ile	Asn	Val	420	425	430
Asp	Trp	Gly	Lys	Pro	Ile	Ile	Asp	Tyr	Ile	Leu	Thr	Gly	Asn	Thr	Ser	435	440	445
Tyr	Pro	Val	Ser	Asp	Asn	Ile	Val	Gln	Val	Asp	Ala	Val	Asp	Gln	Trp	450	455	460
Thr	Tyr	Trp	Leu	Ile	Glu	Asn	Asp	Pro	Glu	Gly	Pro	Phe	Ser	Leu	Pro	465	470	475 480
His	Pro	Met	His	Leu	His	Gly	His	Asp	Phe	Leu	Val	Leu	Gly	Arg	Ser	485	490	495
Pro	Asp	Val	Pro	Ala	Ala	Ser	Gln	Gln	Arg	Phe	Val	Phe	Asp	Pro	Ala	500	505	510
Val	Asp	Leu	Ala	Arg	Leu	Asn	Gly	Asp	Asn	Pro	Pro	Arg	Arg	Asp	Thr	515	520	525

Thr	Met	Leu	Pro	Ala	Gly	Gly	Trp	Leu	Leu	Leu	Ala	Phe	Arg	Thr	Asp
530						535					540				
Asn	Pro	Gly	Ala	Trp	Leu	Phe	His	Cys	His	Ile	Ala	Trp	His	Val	Ser
545					550					555					560
Gly	Gly	Leu	Ser	Val	Asp	Phe	Leu	Glu	Arg	Pro	Ala	Asp	Leu	Arg	Gln
				565					570					575	
Arg	Ile	Ser	Gln	Glu	Asp	Glu	Asp	Asp	Phe	Asn	Arg	Val	Cys	Asp	Glu
			580					585					590		
Trp	Arg	Ala	Tyr	Trp	Pro	Thr	Asn	Pro	Tyr	Pro	Lys	Ile	Asp	Ser	Gly
		595					600					605			
Leu	Lys	Arg	Arg	Arg	Trp	Val	Glu	Glu	Ser	Glu	Trp	Leu	Val	Arg	
610						615					620				

<210> 3
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 3
 tgccacactg cgcccaaccg tgct

24

<210> 4
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 4
 gttcttgata taccaatcag gatg

24

<210> 5
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: primer

<400> 5
 gccggtgagg atgtagtcga tgat

24

<210> 6
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 6

aggtgacgtt gaaccagtag ttgtc

25

<210> 7

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 7

ctggtgcact tcacgcagaa caa

23

<210> 8

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 8

agaaccactt ccaggtgtcg ct

22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 01/00503

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C12N 9/02 // (C12N 9/02, C12R 1:645), D21C 5/00, A61K 7/13, C11D 3/386
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Applied and Environmental Microbiology, Volume 63, No 8, August 1997, Randy M. Berka et al, "Characterization of the Gene Encoding an Extracellular Laccase of Myceliophthora thermophila and Analysis of the Recombinant Enzyme Expressed in Aspergillus oryzae", page 3151 - page 3157, see fig. 4 and abstract --	1-39
X	Can. J. Microbial, Volume 46, 2000, A.M. Farnet et al, "Purification, partial characterization, and reactivity with aromatic compounds of two laccases from Marasmius quercophilus strain 17", page 189 - page 194, see abstract and fig. 6 --	1-39

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

1 November 2001

Date of mailing of the international search report

07-11-2001

Name and mailing address of the ISA/
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Yvonne Siösteen/BS
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/FI 01/00503

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WPI/Derwent's abstract, Accession no. 1997-206626, week 9719, ABSTRACT OF JP,9056378 (DAIWA KASEI KK), 4 March 1997 (4.3.97) --	1-39
X	WPI/Derwent's abstract, Accession no. 1996-203142, week 9621, ABSTRACT OF JP,8070861 (DAIWA KASEI KK) 19 March 1996 (19.03.96) --	1-39
X	US 5750388 A (BERKA ET AL), 12 May 1998 (12.05.98), see fig. 4 and 5 --	1-39
A	EP 0852260 A1 (NOVO NORDISK A/S), 8 July 1998 (08.07.98) --	1-39
A	J.Anim. Morphol. Physiol., Volume 40, No 1 & 2, 1993, M.S. Bharathi et al, "Phenoloxidase associated with the bysall thread tanning in the marine bivalve perna viridis (linnaeus)" page 153 - page 161 --	1-39
A	BIOSIS, PREV199800345876, Vol 27, No 3, page 217-223, Machuca Angela et al: "Production and characterization of thermostable phenol oxidases of the ascomycete Thermoascus aurantiacus", Biotechnology and Applied Biochemistry, June 1998 -- -----	1-39

INTERNATIONAL SEARCH REPORT

Information on patent family members

01/10/01

Internat. application No.

PCT/FI 01/00503

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
US	5750388	A	12/05/98	AT	196164 T	15/09/00
				AU	2656695 A	04/01/96
				DE	69518751 D,T	15/02/01
				EP	0763115 A,B	19/03/97
				SE	0763115 T3	
				ES	2153037 T	16/02/01
				PT	763115 T	30/03/01
				US	5843745 A	01/12/98
				WO	9533837 A	14/12/95

EP	0852260	A1	08/07/98	AU	1192797 A	22/01/97
				JP	11299483 A	02/11/99
				WO	9700948 A	09/01/97
				JP	7302685 A	14/11/95
				JP	11299482 A	02/11/99
